

/// PHASE 2 - CARB CONTROLLER

[What is Phase 2?](#)[Clinical Study](#)[Application](#)[ASIAN Products](#)

NO OTHER WHITE KIDNEY BEAN CAN BEAT PHASE 2®



The Latest, Most Powerful Scientific Evidence of Phase 2®



Article

Systematic Review and Meta-Analysis of a Proprietary Alpha-Amylase Inhibitor from White Bean (*Phaseolus vulgaris* L.) on Weight and Fat Loss in Humans

Jay Udani *, Ollie Tan and Jhanna Molina

Medical Private Practice, Agoura Hills, CA 91301, USA; ollietan124@yahoo.com (O.T.); jhannamolina@gmail.com (J.M.)

* Correspondence: drjayu2016@gmail.com; Tel.: +1-818-885-8500

Received: 14 February 2018; Accepted: 12 April 2018; Published: 20 April 2018



The latest (published Apr 2018) systematic review & meta-analysis on the effectiveness of Phase 2® in weight loss and reducing body fat produced the following findings:


- Phase 2® supplementation showed an average effect on weight loss difference of **-1.08 kg (p less than 0.00001)**
- Phase 2® supplementation showed an average effect on body fat reduction of **3.26 kg (p = 0.02)**

More than two dozens of clinical studies with more than 600 subjects have assessed and confirmed the safety and efficacy of Phase 2® Carb Controller in glycemic control and weight management.

I. PHASE 2[®] RESULTED IN REDUCED POSTPRANDIAL GLUCOSE RESPONSE BY 66% IN 1ST HOUR

Vinson, J. A., Kharrat, H. A., & Shuta, D. (2009). Investigation of an amylase inhibitor on human glucose absorption after starch consumption. *The Open Nutraceuticals Journal*, 2(1).


In a study conducted in University of Scranton (USA), 11 fasting healthy subjects consumed 1500mg of Phase 2 or placebo, together with 4 slices of bread (60g carbohydrate). Subjects' blood glucose level was measured before and after meal periodically every 15 mins up to 2 hours. Phase 2[®] resulted in reduced postprandial glucose response by 66% in 1st hour.



2. EFFICACY OF PHASE 2[®] IN WEIGHT LOSS AND WEIGHT MAINTENANCE

Grube, B., Chong, W. F., Chong, P. W., & Riede, L. (2014). Weight reduction and maintenance with IQP-PV-101: A 12-week randomized controlled study with a 24-week open label period. *Obesity*, 22(3), 645-651.

In the recent most comprehensive human study conducted involving 123 healthy participants, subjects taking a proprietary extract of the white bean, Phase 2 Carb Controller[®] lost an average of seven pounds more than those on placebo after 12 weeks. In addition, 73.5% of the participants in the weight management phase successfully maintained their body weight after 24 weeks.




3. PHASE 2® IMPROVED BODY COMPOSITION OF OVERWEIGHT HUMAN SUBJECTS IN EUROPE

Celleno, Leonardo, et al. "A dietary supplement containing standardized Phaseolus vulgaris extract influences body composition of overweight men and women." *International journal of medical sciences* 4.1 (2007): 45.

A clinical study was conducted by researchers in Celleno University of Rome, Italy on conducted on 60 overweight volunteers. Subjects were given either one 445mg of Phase 2® supplement or a placebo pill daily for 30 consecutively days before a carb-loaded meal. Various body compositions of these subjects were measured after 30 days.

Subjects receiving Phase 2® supplement with a carbohydrate-rich (2000- to 2200-calorie diet) had significantly (p less than 0.001) greater reduction of body weight, BMI, fat mass, adipose tissue thickness, and waist/hip/thigh circumferences while maintaining lean body mass compared to subjects receiving placebo.




4. PHASE 2[®] DECREASES BODY WEIGHT AND WAIST CIRCUMFERENCE OF CHINESE SUBJECTS


Perricone, Nicholas V. "Enhanced weight loss from a dietary supplement containing standardized Phaseolus vulgaris extract in overweight men and women." *METHODS 12.13 (2010): 15-22.*

A randomized, double blinded, and placebo-controlled investigation was carried out on 101 subjects by researchers in Zhejiang University, Hangzhou, China. Subjects were given either 1,000 mg Phase 2[®] or placebo 15 minutes before each meal for 60 consecutive days.


Body weight, waist and hip circumferences, and blood chemistry values were measured at the beginning, at 30 days, and at the end of the 60-day treatment phase. The results showed very significant (p less than 0.001) reduction of body weight, BMI and waist circumference.




PROFESSIONAL ENDORSEMENT BY HEALTHCARE EXPERTS




Steven Rosenblatt, M.D.



Dr. Nicholas Perricone



Sherry Torkos, Pharmacist & Author



Dr. Harry G. Preuss

**MORE THAN 600 MEDIA NEWS COVERAGE WORLDWIDE
SINCE YEAR 2002!**



///PHARMACHEM PACIFIC ASIA

Pharmachem Pacific Asia Pte Ltd, the Asia Representative of Pharmachem Laboratories Inc., USA, has been successfully marketing and distributing health supplements and functional ingredients that aim to improve health and quality of life of consumers through food, beverage and nutritional supplement industries. [> more](#)

Copyright 2018 Pharmachem Pacific Asia
